

## About

Berita Kedokteran Masyarakat (BKM Public Health and Community Medicine) [ISSN [0215-1936](#) (Print) and ISSN [2614-8412](#) (Online)] is a scientific journal in the field of Public Health, published by the Master Program in Public Health, Faculty of Medicine, Public Health, and Nursing (FKKMK) Gadjah Mada University in collaboration with the Association of Indonesian Community Medicine and Public Health (PDK3MI), Minister of Information of the Republic of Indonesia No. 1143 / SK / Ditjen.PPG / STT / 1987 and the [Indonesian Society of Public Health Experts \(IAKMI\)](#).

Berita Kedokteran Masyarakat (BKM Public Health and Community Medicine) is a scientific journal that contains research articles published four months since it was published in 1985, then since 2016 (Volume 32) BKM is published every month / 12 times a year. BKM is recognized as an accredited national scientific journal ([Sinta 3](#)).

Articles published in this journal have gone through a blind review process by two of the reviewer. The reviewer then makes a decision based on a recommendation with several possibilities: rejected, major revision, minor revision, or accepted.

## Make Submission

Focus and Scope

Publication Frequency

Peer Review Process

Open Access Policy

Publication Ethics

Copyright & License

Author Guidelines

Articles Processing Charge

Plagiarism Policy

Indexing

Generative AI

Scopus Citedness

Contact Us

## Download



## MOST READ LAST WEEK

Traffic accidents and injury patterns in underage driving: a study from Klaten Soeradji Tirtonegoro Hospital data  
 359

Determinants of personal hygiene behavior amongst food handlers in fast food restaurants in South Jakarta, Indonesia  
 261

Social media engagement patterns in relation to adolescent anxiety and depression: a systematic review  
 255

Psychosocial factors and work stress in civil servants: a scoping review  
 242

Factors influencing microplastic contamination in bottled drinking water in Indonesia: a systematic review  
 170

## User Account

Username

Password

Remember me

## Statistics

### Visitors

|           |        |
|-----------|--------|
| ID 59,571 | IN 604 |
| SG 5,246  | GB 381 |
| CN 4,030  | JP 342 |
| US 3,782  | MY 304 |
| PH 795    | AU 275 |

Pageviews: 224,711

FLAG counter

**01505719** [View Data](#)

**Editorial Office:** Sekretariat Bersama Jurnal, Lt. 2 Atas Kantin IKM, Jl. Farmako, Sekip Utara, Yogyakarta 55281; Telp: (0274)-547490, Email: [bkm.fk@ugm.ac.id](mailto:bkm.fk@ugm.ac.id)



Berita Kedokteran Masyarakat is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](#).

Platform &  
workflow by  
**OJS / PKP**

## EDITORIAL TEAM

## EDITOR IN CHIEF

Yayi Suryo Prabandari

Universitas Gadjah Mada, Indonesia



## EDITORIAL BOARD


Asnawi Abdullah

Universitas Muhammadiyah Aceh, Indonesia



Bayu Satria Wiratama

Universitas Gadjah Mada, Indonesia



Budi Utomo

Universitas Airlangga, Indonesia



Defriman Djafri

Universitas Andalas, Indonesia



Dewi Susanna

Universitas Indonesia, Indonesia



Dian Mawarni

Universitas Negeri Malang, Indonesia



Mubasysyir Hasanbasri

Universitas Gadjah Mada, Indonesia



Sukri Palutturi

Universitas Hasanuddin, Indonesia



Supriyati

Universitas Gadjah Mada, Indonesia



Annisa Utami Rauf

Universitas Gadjah Mada, Indonesia



## EDITORIAL ASSISTANTS

Vena Jaladara

Universitas Gadjah Mada, Indonesia



Ignasia Ika Kusumaningtyas

Universitas Gadjah Mada, Indonesia



## Make Submission

Focus and Scope

Publication Frequency

Peer Review Process

Open Access Policy

Publication Ethics

Copyright &amp; License

Author Guidelines

Articles Processing Charge

Plagiarism Policy

Indexing

Generative AI

Scopus Citedness

Contact Us

## Download



## MOST READ LAST WEEK

Traffic accidents and injury patterns in underage driving: a study from Klaten Soeradji Tirtonegoro Hospital data  
359

Determinants of personal hygiene behavior amongst food handlers in fast food restaurants in South Jakarta, Indonesia  
261

Social media engagement patterns in relation to adolescent anxiety and depression: a systematic review  
255

Psychosocial factors and work stress in civil servants: a scoping review  
242

Factors influencing microplastic contamination in bottled drinking water in Indonesia: a systematic review  
170

## User Account

Username Password  Remember me

## Statistics

## Visitors

|           |        |
|-----------|--------|
| ID 59,571 | IN 604 |
| SG 5,246  | GB 381 |
| CN 4,030  | JP 342 |
| US 3,782  | MY 304 |
| PH 795    | AU 275 |

Pageviews: 224,711



01505714 View Data

**Editorial Office:** Sekretariat Bersama Jurnal, Lt. 2 Atas Kantin IKM, Jl. Farmako, Sekip Utara, Yogyakarta 55281; Telp: (0274)-547490, Email: bkm.fk@ugm.ac.id



Berika Kedokteran Masyarakat is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).

Platform &  
workflow by  
OJS / PKP

HOME / ARCHIVES / Vol 41 No 09 (2025)


DOI: <https://doi.org/10.22146/bkm.v41i09>

PUBLISHED: 2025-09-16

## ARTICLES

**Social media engagement patterns in relation to adolescent anxiety and depression: a systematic review**


Gorry Amanda, Deby Aprilia Haryani, Yulia Ratna Sofa, Dannu Purwanto, Adiva Kalila Maryam, Ray Wagiu Basrowi, Yuli Puspita Devi e22664



Abstract views: 760 | PDF views: 395

**A scoping review of partners and in-law roles in postpartum depression**


Lintang Kemuning Mustikaning Rahayu, Zahroh Shaluhayah, Antono Suryoputro e23520



Abstract views: 213 | PDF views: 141

**Utilizing "Google Trends" data to support early detection of epidemic outbreaks: a preliminary study**


Frisca Rahmadina, Bagas Suryo Bintoro, Aditya Lia Ramadana e22594



Abstract views: 145 | PDF views: 109

**The time is ripe: a scoping review of risk factors and barriers in preconception care**


Halfie Zaqiyah Gusti Puspitasari, Widyawati, Elsi Dwi Hapsari, Agung Subakti Nuzulullail, Dana Prayoga Irawan e23771



Abstract views: 312 | PDF views: 242

**Origami activities are effective in improving fine motor skills of stunted children**

Uswatun Khasanah, Anindhita Almadevy M.J, Tiar M Pratamawati, Eni Suhaeni, Catur Setiya Sulistiyana, Dea Triasari Indriyanti Wahidin e18366



Abstract views: 183 | PDF views: 152

**Make Submission**

Focus and Scope

Publication Frequency

Peer Review Process

Open Access Policy

Publication Ethics

Copyright &amp; License

Author Guidelines

Articles Processing Charge

Plagiarism Policy

Indexing

Generative AI

Scopus Citedness

Contact Us

**Download****MOST READ LAST WEEK**Traffic accidents and injury patterns in underage driving: a study from Klaten Soeradji Tirtonegoro Hospital data  
 359Determinants of personal hygiene behavior amongst food handlers in fast food restaurants in South Jakarta, Indonesia  
 261Social media engagement patterns in relation to adolescent anxiety and depression: a systematic review  
 255Psychosocial factors and work stress in civil servants: a scoping review  
 242Factors influencing microplastic contamination in bottled drinking water in Indonesia: a systematic review  
 170**User Account**Username Password  Remember me**Statistics****Visitors**

|           |        |
|-----------|--------|
| ID 59,571 | IN 604 |
| SG 5,246  | GB 381 |
| CN 4,030  | JP 342 |
| US 3,782  | MY 304 |
| PH 795    | AU 275 |

Pageviews: 224,711

**01505715** [View Data](#)**Editorial Office:** Sekretariat Bersama Jurnal, Lt. 2 Atas Kantin IKM, Jl. Farmako, Sekip Utara, Yogyakarta 55281; Telp: (0274)-547490, Email: [bkm.fk@ugm.ac.id](mailto:bkm.fk@ugm.ac.id)Berika Kedokteran Masyarakat is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).

## Origami activities are effective in improving the fine motor skills of stunted children

**Submitted:**

December 14th, 2024

**Accepted:**

September 25th, 2025

**Published:**

September 30th, 2025

<sup>1</sup>Department of Public Health,  
Faculty of Medicine,  
Universitas Swadaya Gunung  
Jati, West Java, Indonesia

<sup>2</sup>Department of Genetics,  
Faculty of Medicine,  
Universitas Swadaya Gunung  
Jati, West Java, Indonesia

<sup>3</sup>Department of Medical  
Education, Faculty of  
Medicine, Universitas Swadaya  
Gunung Jati, West Java,  
Indonesia

**\*Correspondence:**

uswatunhasanah7@gmail.com

Uswatun Khasanah<sup>1\*</sup>, Anindhita Almadevy MJ<sup>1</sup>, Tiar M Pratamawati<sup>2</sup>, Eni Suhaeni<sup>1</sup>, Catur Setiya Sulistiyana<sup>3</sup>, Dea Triasari Indriyanti Wahidin<sup>1</sup>

**Abstract**

**Purpose:** This study aimed to evaluate the effectiveness of origami activities on fine motor development in stunted children aged 3–5 years.

**Methods:** A quasi-experimental study with a pretest–posttest control group design was conducted at the Babakan Community Health Center, Cirebon District. A total of 60 stunted children were recruited and equally assigned to intervention and control groups (n = 30 each group). The intervention group participated in three origami activity sessions, while the control group did not receive the intervention. Fine motor skills were assessed before and after the intervention using the KPSP questionnaire. Group differences were analyzed using the Mann–Whitney test. **Results:** Children in the intervention group showed significantly greater improvement in fine motor skills compared with the control group (p = 0.014). **Conclusion:** Origami activities are effective in enhancing fine motor development among stunted children and may serve as a simple, low-cost intervention in community health settings.

**Keywords:** children; fine motor skills; origami; stunting

## INTRODUCTION

Stunting is a condition of impaired growth characterized by a child's height being significantly below the standard for their age. This condition results from chronic malnutrition that often occurs during the critical first 1,000 days of life, from conception to the age of two years. Beyond physical growth retardation, stunting also has profound impacts on brain development, including structural and functional alterations caused by delayed cerebellar neuronal differentiation and shortening of apical dendrites. In Indonesia, stunted growth remains a significant public health challenge [1,2].

The national basic health research (Riset Kesehatan Dasar; Riskesdas) in 2018 reported that 21.6% of Indonesian children are stunted. A nutrition survey, Survei Status Gizi Indonesia (SSGI), in 2022, revealed a stunting prevalence of 20.2% in West Java province. Additionally, stunting incidence in Cirebon city in 2021

was reported to be 17.0% [3,4]. Risk factors for stunting include poor parenting, inadequate health services, limited access to diverse food sources, and inadequate access to clean water and sanitation. Stunting management should involve not only the control of these risk factors to prevent new cases, but also measures to improve the growth and development of stunted preschool children. They also require improvement in both gross and fine motor skills, which are essential for overall development [5-7].

Children aged 1 to 6 years are in a critical period of development during which they are highly responsive to various forms of stimulation. At this stage, they naturally exhibit intense curiosity and eagerness to explore their environment, which makes it essential to provide adequate support for their motor development. Motor development is generally categorized into gross motor skills and fine motor skills. The optimal development of fine motor skills during early childhood is particularly critical, as it lays the

foundation for later learning, independence in daily activities, and overall functional abilities [5,6].

One common issue faced by children with stunting, besides impaired physical growth, is reduced brain function, which affects motor skills, attention, memory, and cognitive abilities. Studies have shown that stunted children experience delays in both gross and fine motor development. These developmental delays limit their ability to be independent and engage in activities, including playing, which can further impact brain, physical, communication, and social development. Therefore, early and adequate interventions are essential to prevent potential losses and support their overall growth and development [5,7-9].

Various strategies have been developed to stimulate fine motor skills in early childhood. Origami, the traditional Japanese art of paper folding, is one such activity that requires hand-eye coordination, concentration, patience, and accuracy. Several studies have demonstrated that origami can enhance fine motor skills, imagination, and spatial abilities in preschool children. Nevertheless, the evidence regarding its effectiveness specifically for stunted children is still limited, particularly in community-based health settings such as posyandu (integrated health posts). This gap highlights the need for empirical studies to investigate the potential of origami as a simple, affordable, and practical intervention for enhancing fine motor development in children with stunted growth in Indonesia [7-10].

Therefore, this study aims to assess the effectiveness of origami activities in improving fine motor skills among stunted children aged 3–5 years in the working area of Babakan Community Health Center, Cirebon. The findings are expected to contribute to the scientific evidence based on developmental stimulation for stunted children and to offer practical implications for incorporating origami-based activities into routine posyandu programs as a complementary strategy to reduce stunting.

## METHODS

This is a quasi-experimental study with a pre- and posttest control group design. The research protocol was approved by the Ethics Committee of the Faculty of Medicine at Universitas Swadaya Gunung Jati (No. 86/EC/FKUGJ/VI/2023). The study population consisted of children aged 3 to 5 years who attended the Posyandu held by the Babakan Community Health Center from June to July 2023.

A total of 60 stunted children were equally assigned to intervention and control groups ( $n = 30$  per group). Due to the limited number of eligible participants,

random allocation was not feasible. To minimize potential bias, matching was conducted based on gender and age when selecting participants for the control group. Family-related information, including maternal education, occupation, and household income, was collected; however, only maternal occupation was used as a matching variable because of time and data constraints. Other potential confounders, such as maternal education, socioeconomic status, and home-based stimulation, were not included in the analysis, which may have influenced the study outcomes.

The sample size was calculated using the standard estimation formula for experimental research, ensuring that the study achieved sufficient statistical power (80%) to detect an actual effect. A standard deviation of 1.195 was applied, and an additional 10% was included to account for potential dropouts. This approach provided adequate sensitivity to minimize both type I and type II errors, thereby strengthening the validity and reliability of the findings. The inclusion criteria were stunted children confirmed through on-site height measurements, while children with congenital disabilities or illnesses at the time of data collection were excluded.

The intervention group participated in origami activities once a week for two hours, for a total of three sessions. Height was measured using a microtoise, and fine motor skills were assessed before and after the intervention using the KPSP questionnaire. The assessors who evaluated fine motor skills were not informed about the children's group assignments or stunting status. However, since stunted children generally have noticeable differences in body size, complete blinding could not be guaranteed. Written informed consent was obtained from the parents of all participating children.

To maintain the consistency of the origami intervention, a structured protocol was created for each session to guide the ordering and type of origami activities provided to participants. Initially, all facilitators had been trained to ensure uniform instruction and interaction. Instruction sessions emphasized uniform procedures, step-by-step demonstrations, and equal practice opportunities for all participants. Furthermore, the research team monitored the implementation of each session to ensure fidelity and minimize variation in intervention delivery.

Data were analyzed using statistical software. Univariate analysis was conducted to describe the characteristics of the study participants and to calculate the mean changes in fine motor skills in both the intervention and control groups. Bivariate analysis

was then performed to assess the effect of origami activities on fine motor skill development among stunted children, using the Mann–Whitney test.

## RESULTS

Table 1 summarizes the demographic characteristics of participants. The intervention and control groups showed comparable distributions in child gender, age, and maternal occupation. Most children were aged 52–60 months, with a predominance of stunting in the intervention group, while all children in the control group were classified as short. Mothers in both groups were mainly housewives, with education levels ranging from elementary to senior high school. Family income was mostly within the Rp 1,510,000–2,000,000 range across both groups.

**Table 1. Characteristics of respondents (n=60)**

| Characteristics of children | Intervention group | Control group   |
|-----------------------------|--------------------|-----------------|
|                             | n = 30<br>n (%)    | n = 30<br>n (%) |
| <b>Gender</b>               |                    |                 |
| Female                      | 15 (50.0)          | 13 (43.3)       |
| Male                        | 15 (50.0)          | 17 (56.7)       |
| <b>Age (months)</b>         |                    |                 |
| 36-41                       | 5 (16.7)           | 6 (20.0)        |
| 42-47                       | 8 (26.7)           | 7 (23.3)        |
| 48-51                       | 3 (10.0)           | 3 (10.0)        |
| 52-60                       | 14 (46.7)          | 14 (46.7)       |
| <b>Stunting category</b>    |                    |                 |
| Short                       | 16 (53.3)          | 30 (100)        |
| Very short                  | 14 (46.7)          | 0 (0)           |
| Characteristics of maternal | Intervention group | Control group   |
|                             | n = 30<br>n (%)    | n = 30<br>n (%) |
| <b>Education</b>            |                    |                 |
| No formal education         | 1 (3.3)            | 0 (0)           |
| Elementary school           | 13 (43.3)          | 3 (10.0)        |
| Junior high school          | 9 (30.0)           | 19 (63.3)       |
| Senior high school          | 7 (23.3)           | 6 (20.0)        |
| Bachelor degree             | 0 (0)              | 2 (6.7)         |
| <b>Occupation</b>           |                    |                 |
| Housewife                   | 23 (76.7)          | 23 (76.7)       |
| Laborer/Farmer              | 2 (6.7)            | 2 (6.7)         |
| Kiosk owner                 | 5 (16.6)           | 5 (16.6)        |
| <b>Family income (Rp)</b>   |                    |                 |
| < 1,000,000                 | 2 (6.7)            | 0 (0)           |
| 1,000,000 – 1,500,000       | 10 (33.3)          | 7 (2.3)         |
| 1,510,000 – 2,000,000       | 15 (50.0)          | 19 (63.3)       |
| > 2,000,000                 | 3 (10.0)           | 4 (13.4)        |

Table 2 shows the mean of changes in fine motor skills among children in each study group. The table indicates that children in the intervention group showed greater improvement in motor skills after the origami activities compared to the control group. This finding suggests that origami activities can enhance fine motor skills among stunted children. In addition, the cadres have been trained to implement origami

activities. This approach aims to enable continued implementation of origami-based activities as part of routine posyandu, thereby supporting long-term behavioral and developmental outcomes in children.

**Table 2. Mean of fine motor skills change**

| Group            | Fine motor skills |                  |                    |
|------------------|-------------------|------------------|--------------------|
|                  | Pre-test mean     | Post-test mean   | Difference         |
| Intervention     | 1.60<br>(SD±0.49) | 2.00<br>(SD±0.0) | 0.400<br>(SD±0.49) |
| Non-intervention | 1.87<br>(SD±0.30) | 2.00<br>(SD±0.0) | 0.133<br>(SD±0.30) |

Note: p (p-value); SD (Standard Deviation)

Table 3 presents the effectiveness of origami activities in enhancing fine motor skills in children with stunted growth, as analyzed using the Mann-Whitney test. Children in the intervention group experienced a significant improvement in fine motor skills after participating in origami activities over three sessions.

**Table 3. The effectiveness of origami activities in improving fine motor skills in stunted children**

| Fine motor skills | Group        |            |           |            | p     |
|-------------------|--------------|------------|-----------|------------|-------|
|                   | Intervention |            | Control   |            |       |
|                   | Pre n (%)    | Post n (%) | Pre n (%) | Post n (%) |       |
| Appropriate       | 18 (60)      | 30 (100)   | 26 (86.7) | 30 (100)   |       |
| Not appropriate   | 12 (40)      | -          | 4 (13.3)  | -          | 0.014 |
| Total             | 30 (100)     | 30 (100)   | 30 (100)  | 30 (100)   |       |

Note: p (p-value)

## DISCUSSION

The most important finding of this study is that origami activities significantly improved fine motor skills among stunted children aged 3–5 years compared with the control group. This demonstrates that even a simple, low-cost, and culturally adaptable intervention can have measurable benefits in stimulating developmental outcomes for children with growth restrictions. For families and communities with limited resources, this kind of intervention is particularly valuable because it is inexpensive, easy to apply, and can be integrated into children’s daily routines [11-15].

Origami can enhance fine motor skills because it requires precise finger movements, grip control, and bimanual coordination, all of which strengthen the small muscles of the hands. At the same time, children must integrate visual perception with motor actions, translating two-dimensional patterns into spatial folds, which supports visual–motor integration and cognitive processing. Repeated practice during origami sessions also facilitates motor learning, while the playful and engaging nature of the activity keeps children

motivated to continue practicing for longer. Prior studies confirm that origami improves dexterity, concentration, and visual-motor integration in pre-schoolers and may serve as a low-cost developmental stimulation for children with stunting, who are at a higher risk of motor delays [5,12,16].

From a practical perspective, these findings carry important implications for community health programs. Integrating origami-based activities into posyandu or early childhood education initiatives can provide an affordable, sustainable, and engaging method to support child development alongside existing nutritional interventions. This approach not only stimulates fine motor skills but also encourages creativity, focus, and social interaction—essential components for school readiness. Policymakers and local health authorities should implement this model into standard programs of child development surveillance and intervention in countries with a high prevalence of stunting [17–18].

The present study highlights the need for future investigations to evaluate the long-term effects of origami interventions and their scalability when implemented by trained community health workers or teachers. Future studies should also integrate more comprehensive family and environmental variables, such as parental education, socioeconomic conditions, and home-based stimulation, to better capture the contextual determinants of developmental outcomes. Additionally, expanding the intervention duration and incorporating follow-up assessments will be essential to determine the sustainability of the observed benefits.

The limitations of this study should be acknowledged. Randomization was not feasible due to the limited number of eligible participants; however, matching was applied based on maternal occupation, gender, and age to minimize group differences. In addition, while assessors were not formally informed about the children's stunting status or group assignment, complete blinding could not be ensured because stunted children are often physically recognizable. Moreover, although maternal occupation was considered for matching, other potential confounders, such as maternal education, household income, and home stimulation, were not fully accounted for, which may have influenced the outcomes. These constraints should be carefully considered when interpreting the findings.

## CONCLUSION

Fine motor skills are crucial for child development, and children with developmental delays often require additional support to foster improvement in this area.

There are various methods to enhance fine motor skills, and origami activities are one such practical approach. In this study, the intervention group showed significant improvement in fine motor skills compared to the control group after three sessions of origami activities. These findings should be integrated into the daily activities of stunted children, including those at school and at home.

Achieving this requires collaboration among the government, parents, and healthcare workers. Further studies are necessary to minimize potential bias by matching the characteristics of subjects in both the intervention and non-intervention groups, as well as by considering a broader range of factors that contribute to stunting. Health institutions, through community-health-center cadres (posyandu), should regularly implement origami activities for children with stunted growth to help address growth and developmental delays.

## REFERENCES

1. Kementerian Kesehatan Republik Indonesia. Buletin stunting. Kementerian Kesehatan RI. 2018. Available from: [\[Website\]](#)
2. Kementerian Kesehatan Republik Indonesia. Pahami dan cegah stunting. Kementerian Kesehatan RI. 2021. Available from: [\[Website\]](#)
3. Badan Penelitian dan Pengembangan Kesehatan KKR. Hasil utama Riskesdas 2018. Jakarta: Kementerian Kesehatan Republik Indonesia; 2018. Available from: [\[Website\]](#)
4. Kementerian Kesehatan Republik Indonesia. Buku saku hasil studi status gizi Indonesia (SSGI) tingkat nasional, provinsi, dan kabupaten/kota tahun 2021. 2021. Available from: [\[Website\]](#)
5. Sulistiyowati, Kartika RI, Sari PIA, Impartina A. The effect of playing origami on the fine motor development among preschool children at Putra Buana Kindergarten in Lamongan Regency. Proceedings of the 2nd International Conference of Health Innovation and Technology (ICHIT 2022). Atlantis Press; 2023. Available from: [\[Website\]](#)
6. Manurung E, Panjaitan RF. Studi komparatif balita stunting di perkotaan dan pedesaan berdasarkan karakteristik keluarga. *Best Journal (Biology Education Science & Technology)*. 2020;3(2):251–6.
7. Khasanah U, Suyatno S, Pradigdo SF. Perbedaan perkembangan antara anak yang stunting dan non-stunting (studi kasus di TK wilayah Pucang Gading, Jawa Tengah, Indonesia). *Jurnal Kesehatan Masyarakat*. 2020;8(2):278–83.
8. Afifah N, Perdani RRW, Utami N. Karakteristik stimulasi ibu dan perkembangan motorik halus

- anak usia 0-3 tahun di Kelurahan Penengahan Raya Kecamatan Kedaton Bandar Lampung. 2018;7(2): 150-4.
9. Putri R, Maghfiroh R, Jumiatmoko J, Hafidah R, Nurjanah NE. Meningkatkan kemampuan motorik halus anak melalui permainan kolase bahan bekas studi literatur. *Jurnal Golden Age*. 2021;5(2):314–22.
  10. Cllaudia ES, Widiastuti AA, Kurniawan M. Origami game for improving fine motor skills for children 4-5 years old in Gang Buaya Village in Salatiga. *Jurnal Obsesi: Jurnal Pendidikan Anak Usia Dini*. 2018;2(2):143-8.
  11. Anisa AN, Syafrudin U, Drupadi R. Playing origami and its impact on fine motor skills development of children aged 4-5. *Journal of Early Childhood Education (JECE)*. 2021;3(1):22–30.
  12. Travers BG, Kirkorian HL, Jiang MJ, Pavalko P. Knowing how to fold ‘em: paper folding across early childhood. *Journal of Motor Learning and Development*. 2021;6(1):147–66.
  13. Kartika C, Suryani Yd, Garna H. Hubungan stunting dengan perkembangan motorik kasar dan halus anak usia 2 – 5 tahun di Desa Panyirapan, Kecamatan Soreang Kabupaten Bandung. *Jurnal Integrasi Kesehatan & Sains*. 2020;2(22).
  14. Wiyono CA, Herliani O, Suhartati S, Indahsari NK, Masfufatun M. Hubungan stunting dengan perkembangan motorik pada anak bawah lima tahun di Puskesmas Kalirungkut Surabaya. *Medika Kartika: Jurnal Kedokteran dan Kesehatan*. 2024; 7(1):12–23.
  15. Murniati C. Perkembangan motorik, bahasa, psikososial balita stunting: literature review. *Jurnal Keluarga Berencana*. 2022;7(1):11–21.
  16. Syahroni A. Pengaruh penggunaan permainan origami terhadap perkembangan motorik halus anak stunting 3-5 tahun: studi di Desa Gunung Batin Udik Kecamatan Terusan Nunyai Kabupaten Lampung Tengah tahun 2020 [thesis]. Semarang: Universitas Diponegoro; 2020. Available from: [Website]
  17. Gonzalez SL, Alvarez V, Nelson EL. Do gross and fine motor skills differentially contribute to language outcomes? A systematic review. *Frontiers in Psychology*. 2019;10:2670.
  18. Nawangsari H, Triuspitasari M, Putrayana M. Pengaruh penggunaan media origami terhadap peningkatan perkembangan motorik halus pada anak usia dini di paud Desa Banjartanggul. *Jurnal Keperawatan*. 2023;21(1):22–30.